

***Cephalophora tropica*: a third European record**

MAŁGORZATA RUSZKIEWICZ-MICHALSKA¹*, PIOTR KNYSAK¹,
IZABELA SKROBEK², ANETA GWIAZDA², SEBASTIAN PISKORSKI² &
JOANNA ŻELAZNA-WIECZOREK¹

¹ Department of Algology and Mycology & ² Myco-Algological Section of the Students Society,
Faculty of Biology and Environmental Protection, University of Łódź,
Banacha 12/16, PL-90-237 Łódź, Poland

* CORRESPONDENCE TO: mrusz@biol.uni.lodz.pl

ABSTRACT—We describe the morphology of *Cephalophora tropica* based on a specimen retrieved from samples of urban water bodies in Łódź city, central Poland, and isolated by a shed snakeskin bait method. We also review the species' distribution to demonstrate its circumglobal range. With previous records from Hungary and Finland, this is only the third report of *C. tropica* from Europe.

KEY WORDS—aquatic environment, conidial fungi, distribution, hyphomycetes, *Pezizales*

Introduction

Cephalophora Thaxt. (*Pezizales*, *Ascomycota*) encompasses hyphomycete species without known teleomorphs (Kirk et al. 2008). Thaxter (1903) originally erected the genus for two coprophilous species: *C. tropica* (type species) and *C. irregularis* Thaxt. [= *C. nigricans* (Bainier) Bainier]. A third species accepted in the genus is *C. uniformis* Narayanan, isolated from jute sacks (Narayanan 1962). Three rotifer- and tardigrade-capturing species were also described in *Cephalophora* as *C. longispora* G.L. Barron et al., *C. muscicola* G.L. Barron et al., and *C. navicularis* Tzean & G.L. Barron (Tzean & Barron 1983, Barron et al. 1990), but it has been suggested that they belong in *Lecophagus* (Dick 1990, Tanabe et al. 1999).

Seifert et al. (2011) refer to *C. tropica* as a pantropical and occasionally temperate species. Our survey of literature citations of *C. tropica* indicates

TABLE 1. Literature review of *Cephalophora tropica* reports

COUNTRY	HOST/SUBSTRATE *	REFERENCE
Africa		
Central African Republic	Dung (<i>Bovidae</i>)	CBS 139.59
Egypt	Air-borne dust	Abdel-Hafez et al. (1990)
		Khallil & Abdel-Sater (1992)
	Moth pupae (<i>Noctuidae</i>)	Ismail & Abdel-Sater (1993)
Ghana	<i>Theobroma</i> (<i>Malvaceae</i>)	Farr & Rossman (2016)
Liberia	Dung (<i>Equidae</i>)	Thaxter (1903)
Asia		
China	Dung (<i>Muridae</i>)	Thaxter (1903)
	Endophytic (<i>Salicaceae</i>)	Zhao & Li (2012)
India	Dung (<i>Cercopithecidae</i>)	Iyer et al. (1973)
	Soil	Barron (1968)
Indonesia	Dung (<i>Muridae</i>)	Thaxter (1903)
Iran	Soil	Pordel et al. (2015)
Israel	<i>Trigonella</i> (<i>Fabaceae</i>)	Chabelska-Frydman (1959)
Japan	Vermicompost	Huang et al. 2013
		Moustafa (1986)
Kuwait	Soil	Al-Musallam (1987, 1990)
		Shahzad & Ghaffar (1995)
Pakistan	<i>Helianthus</i> (<i>Asteraceae</i>)	Sharfun-Nahar & Hashmi (2005)
		Mukhtar (2010)
		Mushtaq & Hashmi (2005)
	<i>Ricinus</i> (<i>Euphorbiaceae</i>)	Dawar et al. (2014)
	Diverse substrates	Ellis (1971)
Russia	Rhizosphere (<i>Sapindaceae</i>)	Zlotnikov et al. (2007)
Thailand	Dung (rodents)	Suwanarit (1986)
Turkey	<i>Capsicum</i> (<i>Solanaceae</i>)	CBS 996.69
Australia & Oceania		
Australia	Diverse substrates	Ellis (1971)
	Painted steel panels; organic materials	Crook & Hindson (1995)
Solomon Is.	Diverse substrates	Ellis (1971)
Europe		
Finland	Wooden parts (sauna)	Salonen & Ruokola (1969)
Hungary	Dung (<i>Cervidae</i>)	Tóth (1994)
		Révay (1998)
North & Central America		
Barbados	Soil	Barron (1968)
Jamaica	Dung (<i>Herpestidae</i>)	Thaxter (1903)
		Minter et al. (2001)
Mexico	<i>Vachellia</i> (<i>Fabaceae</i>)	Ale-Agha et al. (2007)
U.S.A.	<i>Roystonea</i> (<i>Arecaceae</i>)	Farr & Rossman (2016)
	<i>Spathiphyllum</i> (<i>Araceae</i>)	
	<i>Helianthus</i> (<i>Asteraceae</i>)	Roberts et al. (1986)
	<i>Nicotiana</i> (<i>Solanaceae</i>)	Wolf (1949)
South America		
Venezuela	Dung (animals; incl. parrots)	Delgado Avila et al. (2002)

* Plant classification acc. to Stevens (2012)

its circumglobal distribution (TABLE 1), showing the most numerous data originating from Asia with no reports from Arctic and Antarctic. The only European records are from Finland (Salonen & Ruokola 1969) and Hungary (Révay 1998, and literature cited therein).

Materials & methods

In 2015, a study of fungal parasites of planktonic algae was conducted in urban, eutrophic aquatic reservoirs, including the artificial flow pond Młynek on the Olechówka river in Łódź, Central Poland (51°43'32"N 19°31'28"E). The river, which flows mainly within the city limits, also supplies several other artificial flow ponds. The Młynek pond (370 × 30–95 m) is part of an urban park system and in close proximity to a large residential area. The pond is used mainly for recreation, but no swimming is allowed.

On 21 July 2015 a water sample was collected from the shoreline of the pond and held in 125 ml plastic containers. Physical parameters measured included conductivity (EC = 483.1 µS/cm), pH (= 8.2), air temperature (= 25.8°C), and water temperature (= 25.7°C).

The sample was transported in an insulated container to the laboratory of the Department of Algology and Mycology, University of Łódź. A bait method was used in order to isolate fungi in the lab (Batko 1975). Three baits (sesame seeds, shed snakeskin, onion peels) were sterilized with hypochlorite solution and UV irradiation in a laminar-flow cabinet; they were distributed onto sterile Petri dishes together with 20 ml of water sample and incubated for 30 days at 24 ± 2°C. Fungi were identified according to Ellis (1971) with the aid of a Nikon SMZ 745T stereomicroscope and Nikon E-400 light microscope (semi-solid preparations) and photographed using a Nikon DS-F1 camera.

Taxonomy

Cephalophora tropica Thaxt., Bot. Gaz. 35: 158 (1903)

PLATE 1

COLONIES appearing light pink to reddish brown from the colour of conidial mass, effuse. Hyphae superficial, colourless, thin-walled, septate, 5.0–7.5 µm diam. CONIDIOPHORES macronematous, with terminal or rarely lateral ampulla, clavate, straight or slightly flexuous, colourless, smooth, 2–3-celled, 24–30 × 18–20 µm; basal cell trapezoidal, in 3-celled conidiophores the middle cell often slightly sloping. CONIDIOGENOUS CELLS terminal, subsphaerical to clavate, 19–29 × 16–20 µm, bearing up to 3 mature and tens of smaller immature conidia visible as balloon-like outgrowths at the surface of the conidiogenous cell. CONIDIA forming simultaneously, solitary, mostly 2–3-septate, 18–24 × 24–37 µm; cylindrical or slightly clavate, basal cell pointed, arising from a prominent wide (≤2.5 µm diam.) hilum; pale to cinnamon brown, basal and topical cells paler than the central ones, in some mature spores basal cell almost colourless; septa darker, brown, ≤2 µm thick.



PLATE 1. *Cephalophora tropica* (LOD AF-33). A: Mature and immature conidiophores and conidia. B1–3: Conidiophores and conidiogenous cells; black arrow (B1) pointing to young two-celled conidium; white arrow (B2, B3) pointing to the youngest conidia visible as balloon-like outgrowths on the conidiogenous cell. C: mature conidia. Scale bars = 10 µm.

SPECIMEN EXAMINED—POLAND: ŁÓDŹ CITY, Młynek artificial flow pond, shore area, 21 July 2015, leg. P. Knysak (LOD AF-33).

Discussion

The size of the conidiophores and conidia reported by us match the lower limits of measurements given in other reports of the species. In conidia formed in the water-derived culture, the maximum number of septa was four. Similarly, Thaxter (1903) and Wolf (1949) reported 2–5-septate (mainly 3-septate) conidia, although subsequent cultural studies by Mayoral et al. (1964) and Weik & Pappelis (1964) noted conidia with up to 8 and 9 septa.

Although both of the latter studies reported conidia that were deformed or with atypical shapes, we have observed only regular conidial forms.

Beside the two main substrates of *C. tropica* (e.g., dung and soil), the diversity of reported substrata covers a range of living and dead plant and animal tissues (TABLE 1) as well as air-borne dust and a painted steel surface. To our knowledge, we are the first to isolate the species from a sample derived from an aquatic ecosystem. Although Kaushal & Singh (2012) reported *C. tropica* from areas surrounding water bodies in India, their photograph of the fungus does not contain necessary information to confirm the identification as *C. tropica*.

The ability to utilize diverse carbon and nitrogen sources was studied by Mayoral et al. (1964) and Weik & Pappelis (1964). *Cephalophora tropica* grew well using all nine tested nitrogen-containing compounds, but fungal sporulation was enhanced by nitrate-N (Weik & Pappelis 1964). Cellobiose and hexoses (fructose, glucose, galactose, and mannose) proved to be the best carbon sources as indicated by mycelium dry weight and change of pH of cultural media (Mayoral et al. 1964). In our experiment, colonies of *C. tropica* developed only on shed snakeskin, suggesting that that substrate had a positive influence on growth. However, species preference toward keratin is still to be tested.

Acknowledgments

The authors are deeply grateful to reviewers Prof. Maria Dynowska (University of Warmia and Mazury, Olsztyn, Poland), Prof. Harish C. Gujani (St. James School of Medicine, Dutch Caribbean), and MSc. Patrícia Oliveira Fiuza (Universidade Estadual de Feira de Santana, Brazil) for critical and constructive comments on the manuscript. The study was financially supported in frame of the statutory funds of Department of Algology and Mycology, University of Łódź, in the years 2015–16.

Literature cited

- Abdel-Hafez SII, Moubasher AH, Barakat A. 1990. Keratinophilic fungi and other moulds associated with air-dust particles from Egypt. *Folia Microbiologica* (Praha) 35: 311–325. <https://doi.org/10.1007/BF02821283>
- Ale-Agha N, Jensen M, Koeppen CW, Hermes M. 2007. Remarkable microfungi from Oaxaca (Mexico) of *Acacia* species. *Communications in Agricultural and Applied Biological Sciences* 72: 857–866.
- Al-Musallam A. 1987. New and noteworthy fungi from Kuwait. *Journal of the University of Kuwait* 14: 337–250.
- Al-Musallam A. 1990. Distribution of keratinophilic fungi in animal folds in Kuwait. *Mycopathologia* 112(2): 65–70. <https://doi.org/10.1007/bf00436500>
- Barron GL. 1968. The genera of *Hyphomycetes* from soil. Baltimore: The Williams & Wilkins Co. 364 p.

- Barron GL, Morikawa C, Saikawa M. 1990. New *Cephalophora* species capturing rotifers and tardigrades. Canadian Journal of Botany 68: 685–690. <https://doi.org/10.1139/b90-089>
- Batko A. 1975. Zarys hydromykologii [Overview of hydromycology]. PWN, Warszawa.
- Chabelska-Frydman C. 1959. *Cephalophora tropica* Thaxter, a rare hyphomycete isolated from seeds in Israel. Bulletin of the Research Council of Israel, Section D, Botany 7(2): 81–84.
- Crook FM, Hindson WR. 1955. An Australian record of *Cephalophora tropica*. Transactions of the British Mycological Society 38: 218–220. [https://doi.org/10.1016/S0007-1536\(55\)80068-X](https://doi.org/10.1016/S0007-1536(55)80068-X)
- Dawar S, Khalid S, Tariq M. 2014. Seed borne mycoflora of castor bean (*Ricinus communis* L.) from Pakistan. Pakistan Journal of Botany 46: 1485–1488.
- Delgado Avila AE, Piñeiro Chavez AJ, Urdaneta Garcia LM. 2002. Mitosporic coprophilous fungi of Zulia State, Venezuela. Revista Científica, Facultad de Ciencias Veterinarias de la Universidad del Zulia 12: 75–81.
- Dick MW. 1990. The systematic position of *Zoophagus insidians*. Mycological Research 94: 347–354. [https://doi.org/10.1016/S0953-7562\(09\)80360-2](https://doi.org/10.1016/S0953-7562(09)80360-2)
- Ellis MB. 1971. Dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew, Surrey. 608 p.
- Farr DF, Rossman AY. 2016. Fungal Databases, Systematic Mycology and Microbiology Laboratory, ARS, USDA. Retrieved September 12, 2016, from <http://nt.ars-grin.gov/fungaldbatabases/>
- Huang K, Li F, Wei Y, Chen X, Fu X. 2013. Changes of bacterial and fungal community compositions during vermicomposting of vegetable wastes by *Eisenia foetida*. Bioresource Technology 150: 235–241. <https://doi.org/10.1016/j.biortech.2013.10.006>
- Ismail MA, Abdel-Sater MA. 1993. Fungi associated with the Egyptian cotton leafworm *Spodoptera littoralis* Boisduval. Mycopathologia 124: 79–86. <https://doi.org/10.1007/BF01103106>
- Iyer R, Ghosh SK, Sarbhoy AK. 1973. Studies on coprophilous fungi – I. Proceedings of the Indian National Science Academy, Part B, Biological Sciences 39: 199–202.
- Kaushal A, Singh R. 2012. *Cephalophora*: reported first time from surrounding area of water bodies of Bhopal (MP), India. Science Secure Journal of Biotechnology 1: 36–38.
- Khallil AM, Abdel-Sater AM. 1992. Fungi from water, soil and air polluted by the industrial effluents of manqabad superphosphate factory (Assiut, Egypt). Journal of Basic Microbiology 31: 83–100. [https://doi.org/10.1016/0964-8305\(92\)90040-U](https://doi.org/10.1016/0964-8305(92)90040-U)
- Kirk PM, Cannon PF, David JC, Stalpers JA (Eds). 2008. Ainsworth and Bisby's dictionary of the fungi. CABI Bioscience, Wallingford.
- Mayoral BM, Kaplan L, Pappelis AJ. 1964. Carbohydrate utilization by *Cephalophora tropica*. Mycologia 56: 626–630. <https://doi.org/10.2307/3756367>
- Minter DW, Rodríguez Hernández M, Mena Portales J, et al. 2001. Fungi of the Caribbean; an annotated checklist. PDMS Publishing, Isleworth.
- Moustafa AF. 1986. Additional species as new records for the fungal flora of Kuwait. Journal of the University of Kuwait 13: 207–216.
- Mukhtar I. 2010. Sunflower disease and insect pests in Pakistan: a review. African Crop Science Journal 17: 109–118. <https://doi.org/10.4314/acsj.v17i2.54204>
- Mushtaq M, Hashmi MH. 2005. Seed-borne mycoflora of sunflower (*Helianthus annuus* L.). Pakistan Journal of Botany 37: 451–457.
- Narayanan SA. 1962. Some interesting micro-fungi from flood affected substrata in Poona, India. Sydowia 16: 77–84.
- Pordel A, Ahmadpour A, Behnia M, Javan-Nikkhah M. 2015. New records of *Hyphomycetes* fungi from Iran. Mycologia Iranica 2: 69–74.
- Révay A. 1998. Review of the *Hyphomycetes* of Hungary. Studia Biologica Hungarica 27–28: 5–74.

- Roberts RG, Robertson JA, Hanlin RT. 1986. Fungi occurring in the achenes of sunflower (*Helianthus annuus*). Canadian Journal of Botany 64: 1964–1971. <https://doi.org/10.1139/b86-260>
- Salonen A, Ruokola AL. 1969. Mycoflora of the Finnish “sauna” (bath-house). Mycopathologia et Mycologia Applicata 38: 327–336. <https://doi.org/10.1007/BF02052879>
- Seifert K, Morgan-Jones G, Gams W, Kendrick B. 2011. The genera of hyphomycetes. CBS Biodiversity Series 9: 997 p.
- Shahzad S, Ghaffar A. 1995. New records of soilborne root infecting fungi in Pakistan. Pakistan Journal of Botany 27: 209–216.
- Sharfun-Nahar MM, Hashmi MH. 2005. Seed-borne mycoflora of sunflower (*Helianthus annuus* L.). Pakistan Journal of Botany 37: 451–457.
- Stevens PF. 2012. Angiosperm Phylogeny Website. Version 12, July 2012 [and more or less continuously updated since]. www.mobot.org/MOBOT/research/APweb/
- Suwanarit P. 1986. Mycological research in Thailand. Proceedings of the Indian National Science Academy, Part B, Biological Sciences 96: 343–355.
- Tanabe Y, Nagahama T, Saikawa M, Sugiyama J. 1999. Phylogenetic relationship of *Cephalophora* to nematophagous hyphomycetes including taxonomic and nomenclatural emendations of the genus *Lecophagus*. Mycologia 91: 830–835. <https://doi.org/10.2307/3761536>
- Thaxter R. 1903. New or peculiar North American *Hyphomycetes*. III. Botanical Gazette 35: 153–159. <https://doi.org/10.1086/328332>
- Tóth S. 1994. Microscopic fungi of the Pilis and the Visegrad Mts, Hungary. Studia Biologica Hungarica 25: 21–57.
- Tzean SS, Barron GL. 1983. A new predatory hyphomycete capturing bdelloid rotifers in soil. Canadian Journal of Botany 61: 1345–1348. <https://doi.org/10.1139/b83-142>
- Weik KL, Pappelis AJ. 1964. Growth of *Cephalophora tropica* in response to various nitrogen sources. Mycologia 56: 650–655. <https://doi.org/10.2307/3756615>
- Wolf FA. 1949. Two unusual conidial fungi. Mycologia 41: 561–564. <https://doi.org/10.2307/3755076>
- Zhao G-H, Li D-W. 2012. First report of *Cephalophora tropica* on NL 351 poplar in China. Journal of West China Forestry Science 41: 46–52.
- Zlotnikov KM, Kazakova V, Vinokurova NG, Zlotnikova K. 2007. Study of chitin-glucan complexes from the soil micromycete *Cephalophora tropica* D3. Applied Biochemistry and Microbiology 43: 449–452. <https://doi.org/10.1134/S0003683807040163>